

Phyllanthus amarus Schum. & Thonn.

Fam. Phyllanthaceae

Ayurvedic name	Bhui-amalaki, Tamalki
Unani name	Bhui-amla
Hindi name	Bhui Aamla
Trade name	Bhui-amla
Parts used	Whole plant



Phyllanthus amarus

Morphological Characteristics

Phyllanthus amarus is an annual herb 60 to 75 cm tall, quite glabrous. Root is stout and woody. Stems are often branched at base and angular. Leaves are numerous, sub-sessile, distichious, stipulate and paripinnate with small leaflets. The leaflets are oblong, having nerve obscure and base rounded. Root is stout tortuous and woody.

Floral Characteristics

Flowers are very minute, shortly pedicelled numerous and axillary and yellowish in colour. Sepals are 5-6, ovate-oblong outer acute, coriaceous with pale margins; disk in both sexes of glands; male flowers 1-3 pedicelled; female flowers are solitary, larger and erect. Stamens are 3, sessile on a short column didynamous, styles minute, reflexed very short. The fruit is capsule, minute, globose and dehiscent. Seeds are with strong parallel and transverse ribs.

Distribution

The plant is distributed throughout India mainly in tropical and subtropical parts of Country.

Harvest Management

- **Crop Maturity and Harvesting:** The crop matures in 80-90 days when it should be harvested; it has maximum active chemical ingredients at fruiting. However, seeds collection is done after 110-120 days old crop.
- **Post-harvest Management:** Whole plant is pulled manually and shade dried. The dried herb is stored in polythene lined gunny bags at cool, well ventilated godowns.
- **Chemical Constituents:** The herb contains three crystalline lignans including phyllanthine and hypophyllanthine (non-bitter part). In addition, five flavonoids have been identified *viz.* quercetine, astralgin, quercitrin, isoquercitrin and rutin. Four alkaloids have also been separated. The total phyllanthin lignans range between 1 to 1.2% in the dry herb.
- **Yield and Cost of Cultivation:** A yield of 2 to 3 t/ha of dry herb is obtained. Rs. 27,500/- is the cost of cultivation for one hectare.

Therapeutic Uses

The plant is widely used to tone-up sluggish liver and also given in chronic liver condition and jaundice. In Unani medicine, the plant is used in jaundice as deobstruent, diuretic, cooling and astringent. In recent studies, the herb and its root have exhibited antiviral actions on Hepatitis-B.

